

AD-773 679

FAILURE OF METHYLPREDNISOLONE TO PRO-  
TECT LEAD-SENSITIZED RATS AGAINST  
ENDOTOXIN

Robert B. Jones, et al

Naval Medical Research Institute  
Bethesda, Maryland


27 June 1973

DISTRIBUTED BY:

**NTIS**

National Technical Information Service  
U. S. DEPARTMENT OF COMMERCE  
5285 Port Royal Road, Springfield Va. 22151

# DISCLAIMER NOTICE



THIS DOCUMENT IS BEST  
QUALITY AVAILABLE. THE COPY  
FURNISHED TO DTIC CONTAINED  
A SIGNIFICANT NUMBER OF  
PAGES WHICH DO NOT  
REPRODUCE LEGIBLY.

AD-773679

## DOCUMENT CONTROL DATA - R &amp; D

1. SECURITY CLASSIFICATION OF TITLE, BODY OF ABSTRACT AND INDEXING ANNOTATION MUST BE ENTERED WHEN THE OVERALL REPORT IS CLASSIFIED		
2. ORIGINATOR'S ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION
FEDERAL BUREAU OF INVESTIGATION WASHINGTON, D. C. 20014		UNCLASSIFIED
		2b. GROUP
3. TITLE		
EFFECT OF CORTICOSTEROIDS ON PROTECTIVE IMMUNITY TO PROTECT LEAD-SENSITIZED RATS AGAINST		
4. TYPE OF REPORT (Type of report and inclusive dates)		
LARGE PROCESS REPORT		
5. AUTHOR (Last name, middle initial, first name)		
JOHN S. L. WILL, AND LUTZ A. KIESOW		
6. REPORT DATE	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
1973	25	8
8. ORIGINATOR'S REPORT NUMBER(S)	9. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
NR011 .0001 .002 .0007 Report No. 1		
10. DISTRIBUTION STATEMENT		
THIS DOCUMENT HAS BEEN APPROVED FOR PUBLIC RELEASE AND SALE: ITS DISTRIBUTION IS UNLIMITED		
11. DISTRIBUTION NOTES	12. SPONSORING MILITARY ACTIVITY	
Reprinted from: INFECTION AND IMMUNITY 1973, 1: 681-684	BUREAU OF MEDICINE AND SURGERY WASHINGTON, D. C.	
13. ABSTRACT		
Corticosteroids, in doses that protect normal rats against lead poisoning, had no effect in lead-sensitized rats.		

Reproduced by  
**NATIONAL TECHNICAL  
 INFORMATION SERVICE**  
 U.S. Department of Commerce  
 Springfield VA 22151



## Failure of Methylprednisolone to Protect Lead-Sensitized Rats Against Endotoxin

ROBERT B. JONES, JAMES L. WISE, AND LUTZ A. KIESOW

Experimental Medicine Division, Naval Medical Research Institute, Bethesda, Maryland 20014

Received for publication 27 June 1973

Methylprednisolone, in doses that protect normal rats against endotoxin, has no effect in lead-sensitized rats.

The administration of lead acetate ( $\text{PbAc}_2$ ) sensitizes both rats (7) and subhuman primates (4) to minute amounts of bacterial endotoxin. The mechanism of this sensitization has not been defined, although it has been suggested that alterations in either the degradation of endotoxin (8), or in carbohydrate metabolism (3, 4), may be important.

Glucocorticoids are demonstrably effective in protecting normal animals against endotoxin (1, 5). The purpose of the present investigation was to determine if this was also the case in lead-sensitized animals.

Under light ether anesthesia, femoral cut-downs were performed on male Sprague-Dawley rats weighing 180 to 220 g, and an intravenous injection of 20 mg of  $\text{PbAc}_2$  dissolved in 0.5 ml of deionized water given. This was immediately followed by an injection of 0.5 ml of *Serratia marcescens* endotoxin (Difco Laboratories) suspended in 0.15 M NaCl buffered to pH 7.4 with 0.02 M sodium phosphate (PBS), after which 9.5 mg of methylprednisolone (Upjohn Co.) dissolved in 0.5 ml of PBS was also given. Deionized water was used as a control for the  $\text{PbAc}_2$  injections in these experiments. However, other investigators have used sodium acetate with no effect on endotoxin induced mortality (2). PBS was used as a control for the endotoxin and methylprednisolone injections.

The animals were observed for 72 h, although most died within the first 12 h. The mean lethal dose was determined for each group according to the method of Litchfield and Wilcoxon (6). No deaths out of 16 were observed in control animals which received only  $\text{PbAc}_2$ , or only methylprednisolone. Two deaths occurred in the 16 animals which received both  $\text{PbAc}_2$  and methylprednisolone, but no endotoxin.

As can be seen from Table 1, methylprednisolone was quite effective in protecting non-lead-treated rats, a single injection causing a fivefold

TABLE 1. Effect of lead acetate and methylprednisolone on endotoxin lethality in rats

Lead acetate (100 mg/kg <sup>a</sup> )	Methylprednisolone (50 mg/kg <sup>a</sup> )	No. of animals	Endotoxin LD <sub>50</sub> <sup>b</sup> (mg/kg <sup>c</sup> )
-	-	65	16.5 (11.6-23.6) <sup>c</sup>
-	+	64	82.7 (66.4-102.9)
+	-	152	$0.45 \times 10^{-3}$ ( $0.34 \times 10^{-3}$ to $0.58 \times 10^{-3}$ )
+	+	178	$0.38 \times 10^{-3}$ ( $0.31 \times 10^{-3}$ to $0.46 \times 10^{-3}$ )

<sup>a</sup> Body weight.

<sup>b</sup> Abbreviation: LD<sub>50</sub>, mean lethal dose.

<sup>c</sup> Numbers in parenthesis are the 95% confidence limits for each LD<sub>50</sub>.

increase in the mean lethal dose. However, in the lead-treated rats, which were approximately 37,000 times more sensitive to endotoxin, the methylprednisolone was without any effect.

This failure of a potent glucocorticoid to protect lead-sensitized rats against endotoxin suggests that lead may produce important qualitative, as well as quantitative, differences in the response of an animal to endotoxin. Furthermore, it would seem to indicate that under certain circumstances the efficacy of steroids in the treatment of septic shock may be a function of other, seemingly unrelated, factors.

We gratefully acknowledge the excellent technical assistance of Stanley Shapiro.

This work was supported by the Bureau of Medicine and Surgery, Navy Department Subtask MR011.0001.002.0007

### LITERATURE CITED

1. Chedid, L., F. Boyer, and M. Saviard. 1951. Action de la cortisone vis à vis de l'infection expérimentale avec *Salmonella typhi* chez la souris. C. R. Acad. Sci. (Paris) 233:713-716.

2. Filkins, J. P. 1973. Effects of lead acetate on sensitivity to shock, intravascular carbon and endotoxin clearances, and hepatic endotoxin detoxification. *Proc. Soc. Exp. Biol. Med.* **142**:471-475.
3. Filkins, J. P. 1973. Hypoglycemia and depressed hepatic gluconeogenesis during endotoxemia in lead-sensitized rats. *Proc. Soc. Exp. Biol. Med.* **142**:915-918.
4. Holper, K., R. A. Trejo, L. Brettschneider, and N. R. DiLuzio. 1973. Enhancement of endotoxin shock in the lead-sensitized subhuman primate. *Surg. Gynecol. Obstet.* **136**:593-601.
5. Kass, E. H. 1960. Effect of corticosteroids and of hormones of pregnancy on the lethal action of bacterial endotoxin. *Ann. N.Y. Acad. Sci.* **88**:107-115.
6. Litchfield, J. T., Jr., and F. Wilcoxon. 1949. A simplified method of evaluating dose-effect experiments. *J. Pharmacol. Exp. Ther.* **96**:99-113.
7. Selye, H., B. Fuchweber, and L. Bertok. 1966. Effect of lead acetate on the susceptibility of rats to bacterial endotoxins. *J. Bacteriol.* **91**:884-890.
8. Trejo, R. A., and N. R. DiLuzio. 1971. Impaired detoxification as a mechanism of lead acetate induced hypersensitivity to endotoxin. *Proc. Soc. Exp. Biol. Med.* **136**:889-893.